The Indirect Impact of Long-Term Overbrowsing on Insects in the Allegheny National Forest Region of Pennsylvania

Michael J. Chips[1,*], Ellen H. Yerger[2], Arpad Hervanek[1], Tim Nuttle[2,3], Alejandro A. Royo[4], Jonathan N. Pruitt[1], Terrence P. McGlynn[5], Cynthia L. Riggall[5], and Walter P. Carson[1]

Abstract - Overbrowsing has created depauperate plant communities throughout the eastern deciduous forest. We hypothesized these low-diversity plant communities are associated with lower insect diversity. We compared insects inside and outside a 60-year-old deer fenced deer exclosure where plant species richness is 5x higher inside versus outside. We sampled aboveground and litter insects using sweep nets and pitfall traps and identified specimens to family. Aboveground insect abundance, richness, and diversity were up to 50% higher inside the fenced exclosure versus outside. Conversely, litter insect abundance and diversity were consistently higher outside the exclosure. Community composition of aboveground insects differed throughout the summer ($P < 0.05$), but litter insects differed only in late summer. Our results demonstrate that the indirect effects of long-term overbrowsing can reduce aboveground insect diversity and abundance, and change composition even when plant communities are in close proximity.

Introduction

The extirpation of *Canis lupus* L. (Gray Wolf) and *Puma concolor* L. (Cougar) combined with lax deer management have caused *Odocoileus virginianus* (Zimmermann) (White-tailed Deer) to become overabundant throughout the eastern deciduous forest, often creating structurally simple and depauperate understory plant communities (Côté et al. 2004, McCabe and McCabe 1997, Ripple et al. 2010). This scenario is particularly true in the Allegheny National Forest region of Pennsylvania where decades of overbrowsing have reduced understory plant diversity by as much as 50–75% (Banta et al. 2005, Carson et al. 2014, Goetsch et al. 2011, Kain et al. 2011, Rooney and Dress 1997) and caused the formation of recalcitrant understory layers (sensu Royo and Carson 2006) dominated by a few unpalatable species that are inimical to biodiversity recovery (Royo and Carson 2006, Royo et al. 2010).

While the deleterious impact of overbrowsing on forest understories is well known (Côté et al. 2004, Rooney and Waller 2003, Waller 2014), the cascading

[1]Department of Biological Sciences, University of Pittsburgh, A234 Langley Hall, Pittsburgh, PA 15260. [2]Department of Biology, Indiana University of Pennsylvania, 114 Weyandt Hall, Indiana, PA 15705. [3]Ecological Services Division, Civil and Environmental Consultants, Inc., 333 Baldwin Road, Pittsburgh, PA 15205. [4]USDA Forest Service Northern Research Station, Forestry Sciences Lab, PO Box 267, Irvine, PA 16329. [5]Department of Biology, California State University Dominguez Hills, Carson, CA. [*]Corresponding author - mjc119@pitt.edu

Manuscript Editor: Daniel Pavuk

effects that browsing has on arthropod communities are less clear. It is predicted that depauperate understories dominated by browse-tolerant plant species will cascade up trophic levels to lower the abundance and diversity of arthropod groups (Hunter and Price 1992, Siemann et al. 1998, Stewart 2001). This cascade may be driven by losses of both structural complexity and food-resource diversity (Stewart 2001). Alternatively, arthropods may simply track plant abundance and thus be somewhat buffered from declines in host-plant diversity (Siemann 1998, Siemann et al. 1998).

To date, studies within the eastern deciduous forest have focused almost entirely on arthropods in the litter layer and have found that overbrowsing indirectly decreases arthropod abundance with variable effects on diversity (Bressette et al. 2012, Brousseau et al. 2013, Christopher and Cameron 2012, Greenwald et al. 2008, Lessard et al. 2012). Oddly, these studies have overlooked aboveground arthropod communities inhabiting the understory (but see Brousseau et al. 2013) or were hampered by small sample sizes (e.g., Bressette et al. 2012). Studies within other forest types in the southwestern United States (Huffman et al. 2009), on islands in the Pacific Northwest (Allombert et al. 2005), and in forests in Europe (Baines et al. 1994, Suominen et al. 2003) have all found that overbrowsing causes decreases in aboveground arthropod richness. If this is a typical response, overbrowsing could be a serious threat to ecosystem services. Arthropods are principal pollinators, seed dispersers, predators, decomposers, and herbivores, and account for the majority of the animal biomass in ecosystems throughout the world (Handel et al. 1981, Wilson 1987, Wise 1993).

Here, we compared arthropod communities within a 60-year-old deer exclosure to an adjacent reference site in the Allegheny National Forest region of Pennsylvania. This is the oldest known deer fence in the eastern deciduous forest. The understory vegetation inside the fenced plot is composed of a diverse layer of forbs, shrubs, and saplings, while the vegetation directly outside, and in much of this region, is depauperate, structurally simple, and often dominated by a single, highly productive fern species, *Dennstaedtia punctilobula* (Michx.) T. Moore (Hay-Scented Fern; Carson et al. 2014, Goetsch et al. 2011, Kain et al. 2011). We tested the hypothesis that long-term deer overbrowsing indirectly decreases the diversity and richness of insects as well as changes insect community composition.

Field-Site Description

We conducted this study within the Allegheny high plateau region of north-central Pennsylvania on State Game Lands #30 (McKean County, 41°38'N, 78°19'W), which is part of the Hemlock–Northern Hardwood Association (Whitney 1990). Further descriptions of the region's natural history can be found in Bjorkbom and Larson (1977) and Hough and Forbes (1943). Deer have overbrowsed this region, as well as much of Pennsylvania, since the 1930s (Carson et al. 2014, Horsley et al. 2003, Redding 1995).

Methods

We compared insect communities between a 0.4-ha plot surrounded by a well-maintained deer fence (height: 2.1 m, mesh size: 15 cm x 20 cm) versus a nearby reference site (~75 m x 55 m). The fence was erected in the late 1940s and excludes White-tailed Deer, but the mesh is likely large enough to allow in smaller vertebrates such as *Marmota monax* (L.) (Woodchuck; Chips et al. 2014). The exclosure and reference site were exactly the same ones used by Goetsch et al. (2011) and Kain et al. (2011) to evaluate the long-term impact of browsing on the understory and overstory, respectively (see Stout 1998 for further details). The overstory of each site was 60–80 years old and characterized in decreasing order of abundance by *Prunus serotina* Ehrh. (Black Cherry), *Acer saccharum* Marshall (Sugar Maple), *Fagus grandifolia* Ehrh. (American Beech), *Acer rubrum* L. (Red Maple), and *Betula* spp. (birch) (Kain et al. 2011).

A diverse and abundant group of wildflowers and shrubs characterized the understory vegetation inside the fenced plot versus the adjacent reference site (percent plant cover: inside exclosure = 63.42 ± 6.10; reference site = 9.28 ± 1.55, $P < 0.05$; mean species richness: inside exclosure = 6.2; reference site = 1.1, $P < 0.05$; Goetsch et al. 2011). Many of these wildflowers peak in abundance in May prior to canopy closure in late spring. A single species, Hay-scented Fern, dominated the reference site and also dominates understories throughout much of the entire region (Carson et al. 2014, Goetsch et al. 2011, Hill and Silander 2001, Rooney and Dress 1997, Royo et al. 2010). Hay-scented Fern is an unpalatable and fairly shade-tolerant species that peaks in biomass in mid- to late June (Hill and Silander 2001, Horsley and Marquis 1983, Horsley et al. 2003, Rooney and Dress 1997, Royo and Carson 2006).

Insect sampling

We sampled insects on 12 May, 29 June, and 26 August 2011 within the exclosure and in the unfenced reference site 200 m away of the same size, elevation, and slope. We sampled litter insects with pitfall traps and aboveground insects with sweep nets. On each date and throughout each site, we collected 20 sweep-net and 20 pitfall-trap samples. Sweep nets were made of fine mesh with a hoop 30 cm in diameter with a 1-m handle. We swept the top 10–30 cm of all vegetation within reach along 10-m transects starting every 10 m with respect to the 75 m edge of the plot and from random locations with respect to the 55 m edge. All sweep-net transects were at least 3 m from the fence or edge of the reference plot and at least 10 m from each other. All pitfall traps were placed within random locations at least 3 m from the fence and at least 10 m away from each other to minimize edge effects. Pitfall traps were half-liter plastic containers, 8 cm diameter x 10 cm height, filled with 2 cm of 95% ethanol. We placed the rim of the pitfall trap flush with the soil surface and collected them after 24 hours (House and Stinner 1983, Williams 1958). With the help of commonly used identification guides (Borror and White 1970, BugGuide.net, Marshall 2007), we identified all insects larger than 5 mm to family. In addition, identifications were verified at the Carnegie Museum of Natural History.

Statistical analyses

We used univariate repeated-measures analysis of variance (RM-ANOVA; Von Ende 2001) with SAS (PROC GLIMMIX, version 9.3.1; SAS Institute, Inc., Cary, NC) to examine differences in insect abundance, richness, and exponential Shannon diversity ($e^{H'}$) throughout the spring and summer. We used this diversity index because it scales Shannon diversity to a comparable measure of species richness (e.g., number of species; see Jost 2006, 2007). The month of sampling was the repeated measure and the treatment was the deer fence. We treated pitfall and sweep-net sampling techniques as independent measures of arthropod communities and analyzed them separately. To assess differences in arthropod community composition, we performed permutational multivariate analysis of variance (PERMANOVA) via ADONIS using the Vegan Package (R core development team; Anderson 2001). Our experimental treatment (fenced exclosure) was not replicated and our analyses were based upon subsamples (pseudoreplicates sensu Hurlburt 1984), thus our results should be interpreted with appropriate caution (see Oksanen 2001 for a wider discussion on this issue). Nonetheless, because deer have been excluded for 60+ years, this study is quite unique.

Results

We found that decades of overbrowsing indirectly reduced aboveground insect abundance ~40%, richness ~45%, and diversity ~50% throughout spring and summer, but this effect was particularly strong in August (fence x time interaction: $P < 0.001$; Fig. 1A, B, C; Table 1). In addition, overbrowsing shifted aboveground



Figure 1. Mean (± S.E.) abundance, family richness, and diversity ($e^{H'}$) of insects larger than 5 mm collected in the spring and summer of 2011 in the Allegheny National Forest region of north-central Pennsylvania in sweep nets (A–C) and pitfall traps (D–F) inside a 60-year-old fenced exclosure and in a nearby reference site.

insect composition over the entire sampling period, particularly in June (Fig. 2A, B, C; Table 2). This shift was primarily driven by higher counts of elaterid beetles in the reference site (19 vs. 6) and more mirid bugs inside of the fenced exclosure (2 vs. 30) (Appendix 1). In contrast, overbrowsing increased litter insect abundance by ~85%, richness by ~30%, and diversity by ~30% throughout spring and summer (Fig. 1D, E, F; Table 1). However, only in August did community composition of litter insects differ between the 2 sites (Fig. 2D, E, F; Table 2). Specifically, browsing increased the abundance of carabid beetles, chrysomelid beetles, and litter ants particularly in August (Appendix 2).

Discussion

Aboveground insects

We found that overbrowsing caused a reduction in the abundance, richness, and diversity of aboveground insects and also changed the composition of these insect

Table 1. Mixed-model ANOVA results to test for differences in abundance, family richness, and diversity ($e^{H'}$) of insects larger than 5 mm caught in sweep nets and pitfall traps in the spring and summer of 2011 within a 60-year-old fenced exclosure and a nearby reference site in the Allegheny National Forest Region in north-central Pennsylvania. * indicates significant *P*-values. See Figure 1 for all fence x time interactions.

	Abundance			Richness			Diversity ($e^{H'}$)		
	df	*F*	*P*	df	*F*	*P*	df	*F*	*P*
Fence									
Sweep Net	1, 114	8.23	0.0049*	1, 114	14.77	0.0002*	1, 114	16.26	0.0001*
Pitfall Trap	1, 114	10.73	0.0014*	1, 114	4.82	0.0301*	1, 114	3.95	0.0492*
Time									
Sweep Net	2, 114	5.43	0.0056*	2, 114	7.14	0.0012*	2, 114	7.46	0.0009*
Pitfall Trap	2, 114	6.68	0.0018*	2, 114	0.75	0.4768	2, 114	0.79	0.4564
Fence x time									
Sweep Net	2, 114	3.50	0.0334*	2, 114	5.27	0.0065*	2, 114	5.74	0.0042*
Pitfall Trap	2, 114	0.11	0.8965	2, 114	0.21	0.8140	2, 114	0.31	0.7374

Table 2. PERMANOVA results to test for compositional differences of insects larger than 5 mm caught in sweep nets and pitfall traps in the spring and summer of 2011 in a 60-year-old deer fence and an adjacent reference site in the Allegheny National Forest region in north-central Pennsylvania. * indicates significant *P*-values. See Appendices 1 and 2 as well as Figure 2 for a breakdown of insect orders and families.

	Sweep net				Pitfall trap			
	df	R^2	*F*	*P*	df	R^2	*F*	*P*
May	1, 26	0.03	0.65	0.8335	1, 33	0.05	1.66	0.1265
June	1, 36	0.11	4.17	0.0010*	1, 36	0.04	1.40	0.2170
August	1, 34	0.05	1.65	0.0815	1, 31	0.10	3.50	0.0145*
Overall	1, 98	0.03	2.89	0.0005*	1, 102	0.02	1.83	0.1045

communities. These reductions and changes were likely caused by the depauperate nature of the understory vegetation that occurred in the reference site versus the rich understory layer that occurred inside the exclosure. Indeed, the general reduction of herbivorous hemipteran insects (Miridae) drove much of this decline across the sampling period (Appendix 1). Our results may apply broadly because overbrowsing has caused the formation of a depauperate understory plant community throughout the region (Carson et al. 2014) and elsewhere (Royo and Carson 2006, Waller 2014). Our findings are consistent with the impact of ungulate browsers on aboveground insects in lowland rainforests in the Pacific Northwest, European woodlands, boreal forests, and Ponderosa forests (Allombert et al. 2005, Baines et al. 1994, Brousseau et al. 2013, Huffman et al. 2009). While results varied throughout spring and summer (Fig. 1A, B, and C), the overall signal was that long-term deer overabundance caused a decline in the diversity, richness, and abundance of aboveground insects and significantly changed arthropod community composition.

Peaks in arthropod abundance and richness later in the season in the reference site may be attributed to high percent cover of Hay-scented Fern, which reaches peak frond density in June and July followed by senesce in August (Hill and Silander 2001). Though we know very little about insect associations with Hay-scented Fern (Balick et al. 1978), high frond density may create a favorable habitat for understory insects in June (e.g., Elaterid beetles; Fig. 2B, Appendix 1). Arthropod abundance is known to track plant abundance rather than plant diversity



Figure 2. Percent relative abundance in the fenced exclosure (Fence) and reference site at each sampling date in 2011 for both sweep nets in (A) May, (B) June, and (C) August, and pitfall trap samples for (D) May, (E) June, and (F) August. Aboveground insect communities were significantly different throughout the summer, but particularly in June and litter insect communities were significantly different only in August (see Table 2). See Appendices 1 and 2 for the relative abundance of each insect family.

in other community types, particularly grasslands (Siemann 1998, Siemann et al. 1998), and dense host concentrations cause increases in insect herbivore abundance (Long et al. 2003, Root 1973).

Litter insects

Overbrowsing increased litter insect richness, abundance, and diversity throughout the sampling period and led to changes in insect community composition only in late summer. Increases in the absolute abundance of predatory ground beetles as well as ants likely drive these changes. Our findings build upon previous studies in North America and Europe, which documented that the indirect effects of overbrowsing led to increases in the absolute abundance of predatory ground beetles (Brousseau et al. 2013, Melis et al. 2006). The mechanism by which browsing increases the abundance of particular litter insects is unclear, but is likely caused by increased temperatures and lower humidity in the litter layer because of decreased plant cover (Stewart 2001). Specifically, browsers likely create environments favorable to arthropods adapted for high-light, dry environments (Gonzalez-Megias et al. 2004, Melis et al. 2006, Suominem et al. 2003). In contrast, some studies found that overbrowsing causes declines in litter arthropod abundance and diversity via an increase in vulnerability to avian predators and changes in soil nutrients (Bressette et al. 2012, Wardle et al. 2001).

Conclusions

We found that aboveground insect richness, abundance, and diversity were higher and insect species composition different inside a >60-year-old deer exclosure versus a nearby reference site. The sharply contrasting understory plant communities that occurred in each site likely caused these differences. In contrast, we found higher abundance, richness and diversity of litter insects in an area exposed to browsing. Our results, and the results of others, call for large-scale, well-replicated experiments that evaluate not only the impact of overbrowsing on arthropod abundance but also how these may subsequently alter entire food webs in the understory as well as in forest canopies over long time scales. We are aware of only one White-tailed Deer exclosure of this advanced age; therefore deer refugia (e.g., tall boulders) could substitute for very old deer exclosures in this approach (Banta et al. 2005, Chollet et al. 2013, Comisky et al. 2005, Rooney 1997). We suggest that differences in the arthropod communities could cascade further up the food chain and impact the abundance and diversity of higher trophic levels, particularly avian insectivores (e.g., Nuttle et al. 2011).

Acknowledgments

Special thanks to R. Androw, R. Davidson, and J. Rawlins, of the Carnegie Museum of Natural History for aid in insect identification. N. Brouwer, A. Domic, E. Griffin, J. Hua, S.C. Pasquini, J. Slyder, and A. Stoler as well as 4 anonymous reviewers provided helpful comments on the manuscript. Thanks to J. Dzemyan and the Pennsylvania Game Commission for providing access to the study site, and S. Williamson and the 2011 Forest Ecology

and Field Techniques in Ecology and Conservation classes at the University of Pittsburgh Pymatuning Laboratory of Ecology for help with data collection. We apppreciate the support of the Biology Department and the School of Graduate Studies and Research at the Indiana University of Pennsylvania.

Literature Cited

Allombert, S., S.A. Stockton, and J.-L. Martin. 2005. A natural experiment on the impact of overabundant deer on forest invertebrates. Conservation Biology 19:1917–1929.

Anderson, M.J. 2001. A new method for non-parametric multivariate analysis of variance. Austral Ecology 26:32–46.

Baines, D., R.B. Sage, and M.M. Baines. 1994. The implications of Red Deer grazing to ground vegetation and invertebrate communities of Scottish native pinewoods. Journal of Applied Ecology 31:776–783.

Balick, M.J., D.G. Furth, and G.A. Cooper-Driver. 1978. Biochemical and evolutionary aspects of arthropod predation on ferns. Oecologia 35:55–89.

Banta, J.A., A.A. Royo, C. Kirschbaum, and W.P. Carson. 2005. Plant communities growing on boulders in the Allegheny National Forest: Evidence for boulders as refugia from deer and a a bioassay of overbrowsing. Natural Areas Journal 25(2):10–18.

Bjorkbom, J.C., and R.G. Larson. 1977. The Tionesta Scenic and Research Natural Areas. General Technical Report NE-31. United States Department of Agriculture, Forest Service, Northeastern Forest Experimental Station, Upper Darby, PA. 24 pp.

Borror, D.J., and R.E. White. 1970. A Field Guide to Insects of America North of Mexico. Houghton Mifflin Co., New York, NY.

Bressette, J. W., H. Beck, and V. B. Beauchamp. 2012. Beyond the browse line: Complex cascade effects mediated by White-tailed Deer. Oikos 121:1749–1760.

Brousseau, P.-M., C. Hébert, C. Cloutier, and S.D. Côté. 2013. Short-term effects of reduced White-tailed Deer density on insect communities in a strongly overbrowsed boreal forest ecosystem. Biodiversity and Conservation 22:77–92.

Carson, W.P., A.A. Royo and C.J. Peterson. 2014. A pox on our land: A case study of chronic deer overbrowsing throughout the Allegheny National Forest region in Pennsylvania. Pp. 400–411, *In* F.S. Gilliam (Ed.). The Herbaceous Layer in Forests of Eastern North America, Second Edition. Oxford University Press, New York, NY.

Chips, M.J., M.R. Magliocca, B. Hasson, and W.P. Carson. 2014. Quantifying deer and turkey leaf-litter disturbances in the eastern deciduous forest: Have nontrophic effects of consumers been overlooked? Canadian Journal of Forest Research 44:1128–1132.

Chollet, S., C. Baltzinger, L. Ostermann, F. Saint-André, and J.-L. Martin. 2013. Importance for forest plant communities of refuges protecting from deer browsing. forest Ecology and Management 289:470–477.

Christopher, C.C., and G.N. Cameron. 2012. Effects of invasive Amur Honeysuckle (*Lonicera maackii)* and White-Tailed Deer (*Odocoileus virginianus*) on litter-dwelling arthropod communities. American Midland Naturalist 167:256–272.

Comisky, L., A.A. royo, and W.P. Carson. 2005. Deer browsing creates rock refugia gardens on large boulders in the Allegheny National Forest, Pennsylvania. American Midland Naturalist 1543(1):201–206.

Côté, S.D., T.P. Rooney, J.-P. Tremblay, C. Dussault, and D.M. Waller. 2004. Ecological impacts of deer overabundance. Annual Review of Ecology, Evolution, and Systematics 35:113–147.

Goetsch, C., J. Wigg, A.A. Royo, T.E. Ristau, and W.P. Carson. 2011. Chronic overbrowsing and biodiversity collapse in a forest understory in Pennsylvania: Results from a 60-year-old deer exclusion plot. The Journal of the Torrey Botanical Society 138:220–224.

Gonzalez-Megias, A., J.M. Gómez, and F. Sánchez-Pinero. 2004. Effects of ungulates on epigeal arthropods in Sierra Nevada National Park (southeast Spain). Biodiversity and Conservation 13:733–752.

Greenwald, K.R., L.J. Petit, and T.A. Waite. 2008. Indirect effects of a keystone herbivore elevate local animal diversity. Journal of Wildlife Management 72:1318–1321.

Handel, S.N., S.B. Fisch, and G.E. Schatz. 1981. Ants disperse a majority of herbs in a mesic forest community in New York State. Bulletin of the Torrey Botanical Club 108:430–437.

Hill, J.D., and J.A. Silander. 2001. Distribution and dynamics of two ferns: *Dennstaedtia punctilobula* (Dennstaedtiaceae) and *Thelypteris noveboracensis* (Thelypteridaceae) in a Northeast mixed hardwoods–hemlock forest. American Journal of Botany 88:894–902.

Horsley, S.B., and D.A. Marquis. 1983. Interference by weeds and deer with Allegheny hardwood reproduction. Canadian Journal of Forest Research 13:61–69.

Horsley, S.B., S.L. Stout, and D.S. DeCalesta. 2003. White-tailed Deer impact on the vegetation dynamics of a northern hardwood forest. Ecological Applications 13:98–118.

Hough, A.F., and R.D. Forbes. 1943. The ecology and silvics of forests in the high plateaus of Pennsylvania. Ecological Monographs 13:299–320.

House, G.J., and B.R. Stinner. 1983. Arthropods in no-tillage soybean agroecosystems: Community composition and ecosystem interactions. Environmental Management 7:23–28.

Huffman, D.W., D.C. Laughlin, K.M. Pearson, and S. Pandey. 2009. Effects of vertebrate herbivores and shrub characteristics on arthropod assemblages in a northern Arizona forest ecosystem. Forest Ecology and Management 258:616–625.

Hunter, M.D., and P.W. Price. 1992. Playing chutes and ladders: Heterogeneity and the relative roles of bottom-up and top-down forces in natural communities. Ecology 73:724–732.

Hurlbert, S.H. 1984. Pseudoreplication and the design of ecological field experiments. Ecological Monographs 54:187–211.

Jost, L. 2006. Entropy and diversity. Oikos 113:363–375.

Jost, L. 2007. Partitioning diversity into independent alpha and beta components. Ecology 88:2427–2439.

Kain, M., L. Battaglia, A.A. Royo, and W.P. Carson. 2011. Over-browsing in Pennsylvania creates a depauperate forest dominated by an understory tree: Results from a 60-year-old deer exclosure. The Journal of the Torrey Botanical Society 138:322–326.

Lessard, J.-P., W.N. Reynolds, W.A. Bunn, M.A. Genung, M.A. Cregger, E. Felker-Quinn, M.N. Barrios-Garcia, M.L. Stevenson, R.M. Lawton, C.B. Brown, M. Patrick, J.H. Rock, M.A. Jenkins, J.K. Bailey, and J.A. Schweitzer. 2012. Equivalence in the strength of deer herbivory on above- and belowground communities. Basic and Applied Ecology 13:59–66.

Long, Z.T., C.L. Mohler, and W. . Carson. 2003. Extending the resource concentration hypothesis to plant communities: Effects of litter and herbivores. Ecology 84:652–665.

Marshall, S.A. 2007. Insects: Their Natural History and Diversity with a Photographic Guide to Insects of Eastern North America. Firefly Books, Richmond Hill, ON, Canada.

McCabe, T.R. and R.E. McCabe. 1997. Recounting white-tails past. Pp. 11–26, *In* W.J. McShea, H. Underwood, and J.H. Rappole (Eds.). The Science of Overabundance: Deer Ecology and Population Management. Smithsonian Institution Press, Washington, DC.

Melis, C., A. Buset, P.A. Aarrestad, O. Hanssen, E.L. Meisingset, R. Andersen, A. Moksnes, and E. Røskaft. 2006. Impact of Red Deer, *Cervus elaphus*, grazing on Bilberry, *Vaccinium myrtillus*, and composition of ground beetle (Coleoptera, Carabidae) assemblage. Biodiversity and Conservation 15:2049–2059.

Nuttle, T., E.H. Yerger, S.H. Stoleson, and T.E. Ristau. 2011. Legacy of top-down herbivore pressure ricochets back up multiple trophic levels in forest canopies over 30 years. Ecosphere 2:397–409.

Oksanen, L. 2001. Logic of experiments in ecology: Is pseudoreplication a pseudoissue? Oikos 94(1): 27–38.

Redding, J. 1995. History of deer population trends and forest cutting on the Allegheny National Forest. Pp. 214–224, *In* K.W. Gottschalk and S.C.L. Fosbroke (Eds.). Proceedings of the 10th Central Hardwood Forest Conference. USDA Forest Service General Technical Report NE-197. Northeastern Forest Experiment Station, Radnor, PA.

Ripple, W.J., T.P. Rooney, and R.L. Beschta. 2010. Large predators, deer, and trophic cascades in boreal and temperate ecosystems. Pp. 141–161, *In* J. Terborgh and J.A. Estes (Eds.). Trophic Cascades: Predators, Prey, and the Changing Dynamics of Nature, 1st edition. Island Press, Washington, DC.

Rooney, T.P. 1997. Escaping Herbivory: Refuge effects on the morphology and shoot demography of the clonal forest herb *Maianthemum canadense*. Journal of the Torrey Botanical Society 124:280–285.

Rooney, T.P., and W.J. Dress. 1997. Species loss over sixty-six years in the ground-layer vegetation of Heart's Content, an old-growth forest in Pennsylvania, USA. Natural Areas Journal 17:297–305.

Rooney, T.P., and D.M. Waller. 2003. Direct and indirect effects of White-tailed Deer in forest ecosystems. Forest Ecology and Management 181:165–176.

Root, R.B. 1973. Organization of a plant-arthropod association in simple and diverse habitats: The fauna of collards (*Brassica oleracea*). Ecological Monographs 43:95–124.

Royo, A.A., and W.P. Carson. 2006. On the formation of dense understory layers in forests worldwide: Consequences and implications for forest dynamics, biodiversity, and succession. Canadian Journal of Forest Research 36:1345–1362.

Royo, A.A., S.L. Stout, D.S. deCalesta, and T.G. Pierson. 2010. Restoring forest herb communities through landscape-level deer herd reductions: Is recovery limited by legacy effects? Biological Conservation 143:2425–2434

Siemann, E. 1998. Experimental tests of effects of plant productivity and diversity on grassland arthropod diversity. Ecology 79:2057–2070.

Siemann, E., D. Tilman, J. Haarstad, and M. Ritchie. 1998. Experimental tests of the dependence of arthropod diversity on plant diversity. The American Naturalist 152:738–750.

Stewart, A.J.A. 2001. The impact of deer on lowland woodland invertebrates: A review of the evidence and priorities for future research. Forestry 74:259–270.

Stout, S.L. 1998. Deer and forest health. Pennsylvania Forests, Spring 1998:14–16

Suominen, O., J. Niemela, P. Martikainen, P. Niemela, and I. Kojola. 2003. Carabidae and Impact of Reindeer grazing on ground-dwelling Curculionidae assemblages in Lapland. Ecography 26:503–513.

Von Ende, C.N. 2001. Repeated-measures analysis: Growth and other time-dependant measures. Pp. 135–157, *In* S.M. Scheiner and J. Gurevitch (Eds.). Design and Analysis of Ecological Experiments, Second Edition. Oxford University Press, Oxford, UK.

Waller, D.M. 2014. Effects of deer on forest herb layers. Pp. 369–399, *In* F.S. Gilliam (Ed.). The Herbaceous Layer in Forests of Eastern North America, Second Edition. Oxford University Press, New York, NY.

Wardle, D.A., G.M. Barker, G.W. Yeates, K.I. Bonner, and A. Ghani. 2001. Introduced browsing mammals in New Zealand natural forests: Aboveground and belowground consequences. Ecological Monographs 71:587–614.

Whitney, G.G. 1990. The history and status of the hemlock–hardwood forests of the Allegheny Plateau. Journal of Ecology 78:443–458.

Williams, G. 1958. Mechanical time-sorting of pitfall captures. Journal of Animal Ecology 27:27–35.

Wilson, E.O. 1987. Little things that run the world (the importance and conservation of invertebrates). Conservation Biology 1:344–346.

Wise, D.H. 1993. Spiders in Ecological Webs. Cambridge University Press, Cambridge, UK.

Appendix 1. Mean relative abundance (± S.E.) and total count of insects larger than 5 mm caught in sweep nets in the spring and summer of 2011 in a 60-year-old deer fence and adjacent reference site in the Allegheny National Forest Region in northcentral Pennsylvania. All insects were identified to family.

		Sweep-net samples		
Order/family	Treatment	May	June	August
Coleoptera				
Canthandae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Cerembycidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Chrysomelidae	Reference	0.11 (0.08)	0.01 (0.01)	0.00 (0.00)
	Fence	0.07 (0.04)	0.01 (0.01)	0.01 (0.01)
Curculionidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
Elateridae	Reference	0.00 (0.00)	0.23 (0.05)	0.00 (0.00)
	Fence	0.02 (0.02)	0.01 (0.04)	0.00 (0.00)
Lucanidae	Reference	0.11 (0.08)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Staphylinidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.02 (0.02)	0.01 (0.01)	0.01 (0.01)
Diptera				
Anthomyiidae	Reference	0.00 (0.00)	0.00 (0.00)	0.11 (0.06)
	Fence	0.00 (0.00)	0.01 (0.01)	0.05 (0.02)
Anthomyzidae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Calliphoridae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
Chironomidae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Dolichopodidae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.02 (0.02)	0.00 (0.00)	0.05 (0.03)
Drosophilidae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.05 (0.03)	0.00 (0.00)	0.00 (0.00)
Empididae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.05 (0.03)	0.00 (0.00)	0.00 (0.00)
Heleomyzidae	Reference	0.06 (0.06)	0.00 (0.00)	0.04 (0.04)
	Fence	0.00 (0.00)	0.03 (0.02)	0.02 (0.01)
Lauxaniidae	Reference	0.00 (0.00)	0.18 (0.09)	0.11 (0.08)
	Fence	0.00 (0.00)	0.05 (0.02)	0.08 (0.03)
Lonchaeidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
Micropezidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
Muscidae	Reference	0.00 (0.00)	0.04 (0.04)	0.04 (0.04)
	Fence	0.00 (0.00)	0.04 (0.02)	0.04 (0.01)
Mycetophilidae	Reference	0.00 (0.00)	0.01 (0.01)	0.04 (0.04)
	Fence	0.00 (0.00)	0.04 (0.02)	0.00 (0.00)
Phoridae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.07 (0.04)	0.00 (0.00)	0.00 (0.00)
Rhagionidae	Reference	0.06 (0.06)	0.00 (0.00)	0.00 (0.00)
	Fence	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)

Order/family	Treatment	Sweep-net samples May	June	August
Simuliidae	Reference	0.11 (0.08)	0.00 (0.00)	0.00 (0.00)
	Fence	0.05 (0.03)	0.00 (0.00)	0.00 (0.00)
Stratiomyidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Syrphidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.01 (0.01)	0.02 (0.02)
Tachinidae	Reference	0.00 (0.00)	0.00 (0.00)	0.04 (0.04)
	Fence	0.07 (0.04)	0.00 (0.00)	0.00 (0.00)
Tipulidae	Reference	0.00 (0.00)	0.05 (0.03)	0.00 (0.00)
	Fence	0.05 (0.03)	0.12 (0.04)	0.01 (0.01)
Hemiptera				
Aphididae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
Berytidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
Cercopidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
Cicadellidae	Reference	0.11 (0.08)	0.01 (0.01)	0.14 (0.07)
	Fence	0.07 (0.04)	0.03 (0.02)	0.09 (0.02)
Membracidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.02 (0.01)	0.04 (0.02)
Miridae	Reference	0.00 (0.00)	0.02 (0.02)	0.11 (0.06)
Nabidae	Reference	0.00 (0.00)	0.02 (0.02)	0.14 (0.08)
	Fence	0.02 (0.02)	0.00 (0.00)	0.16 (0.05)
Pentatomidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.04 (0.01)
Hymenoptera				
Braconidae	Reference	0.00 (0.00)	0.00 (0.00)	0.04 (0.04)
	Fence	0.02 (0.02)	0.01 (0.01)	0.01 (0.01)
Formicidae	Reference	0.11 (0.08)	0.00 (0.00)	0.00 (0.00)
	Fence	0.10 (0.04)	0.00 (0.00)	0.00 (0.00)
Ichneumonidae	Reference	0.00 (0.00)	0.05 (0.02)	0.14 (0.07)
	Fence	0.00 (0.00)	0.03 (0.02)	0.09 (0.03)
Pteromalidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
Tenthredinidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.05 (0.03)	0.05 (0.04)	0.05 (0.02)
Lepidoptera				
Arctiidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Galechiidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
Geometridae	Reference	0.06 (0.06)	0.06 (0.02)	0.04 (0.04)
	Fence	0.02 (0.02)	0.00 (0.00)	0.01 (0.01)
Noctuidae	Reference	0.00 (0.00)	0.04 (0.02)	0.00 (0.00)
	Fence	0.00 (0.00)	0.02 (0.01)	0.02 (0.01)
Notodontidae	Reference	0.00 (0.00)	0.04 (0.02)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.02 (0.02)
Mecoptera				
Panorpidae	Reference	0.00 (0.00)	0.15 (0.04)	0.00 (0.00)
	Fence	0.00 (0.00)	0.18 (0.05)	0.00 (0.00)

Order/family	Treatment	Sweep-net samples		
		May	June	August
Neuroptera				
Hemerobiidae	Reference	0.00 (0.00)	0.00 (0.00)	0.04 (0.04)
	Fence	0.05 (0.05)	0.00 (0.00)	0.00 (0.00)
Orthoptera				
Acrididae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
Psocodea				
Pseudocaecilidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Total count	Reference	18	82	28
	Fence	42	106	141

Appendix 2. Mean relative abundance (± S.E.) and total count of insects larger than 5 mm caught in pitfall traps in the spring and summer of 2011 in a 60-year-old deer fence and adjacent reference site in the Allegheny National Forest Region in north-central Pennsylvania. All insects were identified to family.

		Pitfall-trap samples		
Order/family	Treatment	May	June	Aug
Coleoptera				
Anthribidae	Reference	0.03 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.03 (0.03)	0.00 (0.00)	0.00 (0.00)
Carabidae	Reference	0.17 (0.10)	0.05 (0.02)	0.35 (0.08)
	Fence	0.24 (0.07)	0.10 (0.04)	0.11 (0.05)
Chrysomelidae	Reference	0.16 (0.05)	0.00 (0.00)	0.05 (0.02)
	Fence	0.11 (0.05)	0.00 (0.00)	0.00 (0.00)
Curculionidae	Reference	0.00 (0.00)	0.01 (0.01)	0.03 (0.03)
	Fence	0.03 (0.03)	0.00 (0.00)	0.00 (0.00)
Elateridae	Reference	0.06 (0.05)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Erotylidae	Reference	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.05 (0.04)	0.00 (0.00)	0.00 (0.00)
Histeridae	Reference	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Lampyridae	Reference	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Nitidulidae	Reference	0.09 (0.05)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Scarabacidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Silphidae	Reference	0.06 (0.04)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Staphylinidae	Reference	0.20 (0.07)	0.14 (0.05)	0.01 (0.01)
	Fence	0.24 (0.09)	0.15 (0.07)	0.08 (0.04)
Tenebrionidae	Reference	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Collembola				
Tomoceridae	Reference	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
	Fence	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
Diptera				
Anisopodidae	Reference	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Anthomyiidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.03 (0.03)
Calliphoridae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.08 (0.04)
Lauxaniidae	Reference	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Muscidae	Reference	0.00 (0.00)	0.01 (0.01)	0.01 (0.01)
	Fence	0.00 (0.00)	0.03 (0.02)	0.03 (0.03)
Nematocera	Reference	0.02 (0.02)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Tipulidae	Reference	0.00 (0.00)	0.06 (0.03)	0.01 (0.01)
	Fence	0.00 (0.00)	0.12 (0.05)	0.00 (0.00)

		Pitfall-trap samples		
Order/family	Treatment	May	June	Aug
Hemiptera				
Cicadellidae	Reference	0.02 (0.02)	0.01 (0.01)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Pentatomidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.03 (0.03)	0.00 (0.00)	0.00 (0.00)
Hymenoptera				
Formicidae	Reference	0.13 (0.05)	0.69 (0.18)	0.48 (0.10)
	Fence	0.29 (0.11)	0.59 (0.2)	0.59 (0.15)
Lepidoptera				
Geometridae	Reference	0.00 (0.00)	0.02 (0.02)	0.01 (0.01)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Noctuidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.05 (0.04)
Mecoptera				
Panorpidae	Reference	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
	Fence	0.00 (0.00)	0.00 (0.00)	0.03 (0.03)
Orthoptera				
Gryllaeridae	Reference	0.00 (0.00)	0.00 (0.00)	0.01 (0.01)
	Fence	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
Total count	Reference	64	116	75
	Fence	38	68	37